
Maryland MCAP Grade 8

Math Made Ridiculously SIMPLE



- Smarter Practice. Higher Scores.
- Fast, Fun, Focused Learning!
- Your Shortcut to Math Success!
- Conquer the Math Test!


A Comprehensive 8th Grade Math Study Guide to Acing the Test

Learn Faster · Understand Deeper · Succeed with Confidence

Maryland MCAP Grade 8 Math

Made Ridiculously Simple

A Comprehensive 8th Grade Math Study Guide to Acing the MCAP Test



Reza Nazari

Copyrighted Material

Copyright © 2026

Effortless Math Education Inc

Published by Effortless Math Education

www.effortlessmath.com

All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the author, except in the case of brief quotations embodied in critical reviews and certain other noncommercial uses permitted by copyright law, including Section 107 or 108 of the 1976 United States Copyright Act.

This publication is independently produced and has no official connection to any state, district, or national testing program.

Test names and organizational names used herein are the property of their respective trademark holders.

Copyright © 2026

Copyrighted Material

8.G | Geometry

Understand congruence, similarity, and apply the Pythagorean Theorem

8.G.A	Understand congruence and similarity using physical models, transparencies, or geometry software
8.G.A.1	Verify experimentally the properties of rotations, reflections, and translations:
8.G.A.2	Understand that a two-dimensional figure is congruent to another if the second can be obtained from the first by a sequence of rotations, reflections, and translations; given two congruent figures, describe a sequence that exhibits the congruence between them
8.G.A.3	Describe the effect of dilations, translations, rotations, and reflections on two-dimensional figures using coordinates
8.G.A.4	Understand that a two-dimensional figure is similar to another if the second can be obtained from the first by a sequence of rotations, reflections, translations, and dilations; given two similar two-dimensional figures, describe a sequence that exhibits the similarity between them
8.G.A.5	Use informal arguments to establish facts about the angle sum and exterior angle of triangles, about the angles created when parallel lines are cut by a transversal, and the angle-angle criterion for similarity of triangles
8.G.B	Understand and apply the Pythagorean Theorem
8.G.B.6	Explain a proof of the Pythagorean Theorem and its converse
8.G.B.7	Apply the Pythagorean Theorem to determine unknown side lengths in right triangles in real-world and mathematical problems in two and three dimensions
8.G.B.8	Apply the Pythagorean Theorem to find the distance between two points in a coordinate system
8.G.C	Solve real-world and mathematical problems involving volume of cylinders, cones, and spheres
8.G.C.9	Know the formulas for the volumes of cones, cylinders, and spheres and use them to solve real-world and mathematical problems

8.NS | The Number System

Understand and apply concepts of rational and irrational numbers

Scan Me

8.NS.A	Know that there are numbers that are not rational, and approximate them by rational numbers
8.NS.A.1	Know that numbers that are not rational are called irrational. Understand informally that every number has a decimal expansion; for rational numbers show that the decimal expansion repeats eventually, and convert a decimal expansion which repeats eventually into a rational number
8.NS.A.2	Use rational approximations of irrational numbers to compare the size of irrational numbers, locate them approximately on a number line diagram, and estimate the value of expressions (e.g., Π2)

8.SP | Statistics and Probability

Investigate patterns of association in bivariate data

8.SP.A	Investigate patterns of association in bivariate data
8.SP.A.1	Construct and interpret scatter plots for bivariate measurement data to investigate patterns of association between two quantities. Describe patterns such as clustering, outliers, positive or negative association, linear association, and nonlinear association
8.SP.A.2	Know that straight lines are widely used to model relationships between two quantitative variables. For scatter plots that suggest a linear association, informally fit a straight line, and informally assess the model fit by judging the closeness of the data points to the line
8.SP.A.3	Use the equation of a linear model to solve problems in the context of bivariate measurement data, interpreting the slope and intercept
8.SP.A.4	Understand that patterns of association can also be seen in bivariate categorical data by displaying frequencies and relative frequencies in a two-way table. Construct and interpret a two-way table summarizing data on two categorical variables collected from the same subjects. Use relative frequencies calculated for rows or columns to describe possible association between the two variables

Table of Contents

★ 1 Real Numbers & Number Sense 13

★ 2 Exponents, Roots, & Scientific Notation 27

★ 3 Proportional Relationships & Slope 42

★ 4 Equations & Systems 54

★ 5 Functions: Foundations 67

★ 6 Functions: Models And Patterns 79

★ 7 Transformations & Similarity 91

★ 8 Geometry & Measurement 106

★ 9 Statistics & Data Modeling 131

★ Grade 8 Math Overview 148

★ Test-Taking Strategies 150

★ Practice Test 1 154


★ Practice Test 2 173

★ Chapter Practices Answers and Explanations 190

1.3 Estimating Irrational Numbers

8.NS.A, 8.NS.A.2

When a square root isn't a perfect square, you can't just write down the exact decimal, but here's the good news: you can estimate it super well! The trick is to squeeze it between two perfect squares. For example, since $36 < 45 < 49$, you immediately know that $6 < \sqrt{45} < 7$—your answer is somewhere in that range. Then test tenths or hundredths to narrow it down even more. **The why behind this:** when you square nearby numbers, they tell you if you're too low or too high. This clever method helps you place irrational numbers on a number line and compare them to regular whole numbers and decimals. You've got a superpower now!



Key Concepts & Quick Review

Squeeze between consecutive integers: Find n and $n+1$ such that $n^2 < x < (n+1)^2$; then $n < \sqrt{x} < n+1$.

Refine to tenths: Try $n.1$, $n.2, \ldots$ until you find $a^2 < x < b^2$; then $a < \sqrt{x} < b$.

Comparing irrationals: $\sqrt{a} < \sqrt{b}$ whenever $a < b$ $(a, b > 0)$, so larger radicand $\Rightarrow$ larger square root.

Examples

① Between which two consecutive integers does $\sqrt{20}$ lie?

Find two perfect squares that squeeze 20: We have $4^2 = 16 < 20$ and $5^2 = 25 > 20$. So $\sqrt{20}$ must be between 4 and 5—you can be sure of that! **Now here's a helpful hint:** since 20 is closer to 25 than to 16, the square root is closer to 5. In fact, $\sqrt{20} \approx 4.47$—pretty close to 5, right?

Answer: $4 < \sqrt{20} < 5$

② Estimate $\sqrt{45}$ to one decimal place.

👉 First, squeeze with integers: $6^2 = 36 < 45 < 49 = 7^2$, so $6 < \sqrt{45} < 7$. Now refine to tenths. **Test:** $6.7^2 = 44.89 < 45$ and $6.8^2 = 46.24 > 45$. This tells us $6.7 < \sqrt{45} < 6.8$—we're narrowing it down! Since 45 is much closer to 44.89, we know $\sqrt{45} \approx 6.7$ is our best estimate to one decimal place.

💡 **Answer:** $\sqrt{45} \approx 6.7$

✎ Practice Problems

Give the two consecutive integers that bound each square root, then estimate to one decimal place where indicated.

1. $\sqrt{10}$ is between _____

2. $\sqrt{18}$ is between _____

3. $\sqrt{35}$ is between _____

4. $\sqrt{50}$ is between _____

5. $\sqrt{72}$ is between _____

6. $\sqrt{90}$ is between _____

7. $\sqrt{6} \approx$ _____ (1 d.p.)

8. $\sqrt{14} \approx$ _____ (1 d.p.)

9. $\sqrt{30} \approx$ _____ (1 d.p.)

10. $\sqrt{52} \approx$ _____ (1 d.p.)

11. Order: $\sqrt{5}$, 2, $\sqrt{3}$ _____

12. Greater: $\sqrt{11}$ or 3.3? _____

Study Tips

- Memorize perfect squares through 169 (13^2). This gives you the power to quickly bracket any square root and know which two integers it falls between. You're building your math toolkit!
- When estimating $\sqrt{x}$ to one decimal place, find the two tenths it falls between, then decide which is closer. If they're equally close, test hundredths for extra accuracy before rounding.
- **Here's a time-saver for comparing irrational numbers:** just compare the radicands directly—no estimation needed! Since $47 < 53$, you know right away that $\sqrt{47} < \sqrt{53}$. Smart thinking!

Scan Me

✍ Word Problems

13. A square garden has an area of 75 m^2. The gardener wants to know whether the side length is closer to 8 m or 9 m so she can decide how many fence posts to order. Between which two consecutive integers does the side length fall, and which integer is it closest to?

_____

14. Two drones take off from the same point and fly in perpendicular directions. Drone A travels $\sqrt{60}$ km and Drone B travels 7.5 km. Without using a calculator, determine which drone flies farther.

_____

15. A park ranger estimates the distance across a circular lake. The lake's area is 200 m^2. (a) The radius satisfies $\pi r^2 = 200$. Find r^2. (b) Estimate r to the nearest tenth. (c) Estimate the diameter to the nearest metre.

_____

16. Between which two consecutive whole numbers does each value lie? (a) $\sqrt{50}$ (b) $\sqrt[3]{100}$ (c) Order from least to greatest: $\sqrt{50}$, $\sqrt[3]{100}$, 7.2

_____

17. A square garden has area 150 ft^2. You need to buy fencing for all four sides. Fencing is sold in whole-foot lengths. (a) Find the exact side length. (b) Estimate the perimeter to the nearest tenth of a foot. (c) How many whole feet of fencing must you buy?

_____