
North Carolina

EOG Grade 8

Math Made Ridiculously SIMPLE



- Smarter Practice. Higher Scores.
- Fast, Fun, Focused Learning!
- Your Shortcut to Math Success!
- Conquer the Math Test!


A Comprehensive 8th Grade Math Study Guide to Acing the Test

Learn Faster · Understand Deeper · Succeed with Confidence

North Carolina EOG Grade 8 Math

Made Ridiculously Simple

A Comprehensive 8th Grade Math Study Guide to Acing the EOG Test



Reza Nazari

Copyrighted Material

Copyright © 2026

Effortless Math Education Inc

Published by Effortless Math Education

www.effortlessmath.com

All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the author, except in the case of brief quotations embodied in critical reviews and certain other noncommercial uses permitted by copyright law, including Section 107 or 108 of the 1976 United States Copyright Act.

This publication is independently produced and has no official connection to any state, district, or national testing program.

Test names and organizational names used herein are the property of their respective trademark holders.

Copyright © 2026

Copyrighted Material

NC.8.SP.2	Model the relationship between bivariate quantitative data to:
NC.8.SP.3	Use the equation of a linear model to solve problems in the context of bivariate quantitative data, interpreting the slope and y-intercept
NC.8.SP.4	Understand that patterns of association can also be seen in bivariate categorical data by displaying frequencies and relative frequencies in a two-way table

Table of Contents

★ 1 Real Numbers & Number Sense 12

★ 2 Exponents, Roots, & Scientific Notation 26

★ 3 Proportional Relationships & Slope 41

★ 4 Equations & Systems 53

★ 5 Functions: Foundations 69

★ 6 Functions: Models And Patterns 81

★ 7 Transformations & Similarity 93

★ 8 Geometry & Measurement 108

★ 9 Statistics & Data Modeling 128

★ Grade 8 Math Overview 145

★ Test-Taking Strategies 147

★ Practice Test 1 151

★ Practice Test 2 170

★ Chapter Practices Answers and Explanations 189

CHAPTER

1

Real Numbers & Number Sense

1. Rational and Irrational Numbers NC.8.NS, NC.8.NS.1
2. Turning Repeating Decimals into Fractions NC.8.NS, NC.8.NS.1
3. Estimating Irrational Numbers NC.8.NS, NC.8.NS.2
4. Estimating Expressions with Irrational Numbers NC.8.NS, NC.8.NS.1

② Estimate $\sqrt{45}$ to one decimal place.

First, squeeze with integers: $6^2 = 36 < 45 < 49 = 7^2$, so $6 < \sqrt{45} < 7$. Now refine to tenths. **Test:** $6.7^2 = 44.89 < 45$ and $6.8^2 = 46.24 > 45$. This tells us $6.7 < \sqrt{45} < 6.8$—we're narrowing it down! Since 45 is much closer to 44.89, we know $\sqrt{45} \approx 6.7$ is our best estimate to one decimal place.

Answer: $\sqrt{45} \approx 6.7$

Practice Problems

Give the two consecutive integers that bound each square root, then estimate to one decimal place where indicated.

1. $\sqrt{10}$ is between _____

2. $\sqrt{18}$ is between _____

3. $\sqrt{35}$ is between _____

4. $\sqrt{50}$ is between _____

5. $\sqrt{72}$ is between _____

6. $\sqrt{90}$ is between _____

7. $\sqrt{6} \approx$ _____ (1 d.p.)

8. $\sqrt{14} \approx$ _____ (1 d.p.)

9. $\sqrt{30} \approx$ _____ (1 d.p.)

10. $\sqrt{52} \approx$ _____ (1 d.p.)

11. Order: $\sqrt{5}$, 2, $\sqrt{3}$ _____

12. Greater: $\sqrt{11}$ or 3.3? _____

Study Tips

- Memorize perfect squares through 169 (13^2). This gives you the power to quickly bracket any square root and know which two integers it falls between. You're building your math toolkit!
- When estimating $\sqrt{x}$ to one decimal place, find the two tenths it falls between, then decide which is closer. If they're equally close, test hundredths for extra accuracy before rounding.
- **Here's a time-saver for comparing irrational numbers:** just compare the radicands directly—no estimation needed! Since $47 < 53$, you know right away that $\sqrt{47} < \sqrt{53}$. Smart thinking!

✍ Word Problems

13. A square garden has an area of 75 m^2. The gardener wants to know whether the side length is closer to 8 m or 9 m so she can decide how many fence posts to order. Between which two consecutive integers does the side length fall, and which integer is it closest to?

_____

14. Two drones take off from the same point and fly in perpendicular directions. Drone A travels $\sqrt{60}$ km and Drone B travels 7.5 km. Without using a calculator, determine which drone flies farther.

_____

15. A park ranger estimates the distance across a circular lake. The lake's area is 200 m^2. (a) The radius satisfies $\pi r^2 = 200$. Find r^2. (b) Estimate r to the nearest tenth. (c) Estimate the diameter to the nearest metre.

_____

16. Between which two consecutive whole numbers does each value lie? (a) $\sqrt{50}$ (b) $\sqrt[3]{100}$ (c) Order from least to greatest: $\sqrt{50}$, $\sqrt[3]{100}$, 7.2

_____

17. A square garden has area 150 ft^2. You need to buy fencing for all four sides. Fencing is sold in whole-foot lengths. (a) Find the exact side length. (b) Estimate the perimeter to the nearest tenth of a foot. (c) How many whole feet of fencing must you buy?


_____

Scan Me

1.4 Estimating Expressions with Irrational Numbers

NC.8.NS, NC.8.NS.1

Estimating a *single* irrational number like $\sqrt{45}$ is one thing—you squeeze it between two perfect squares and zero in on the decimal. But what happens when an expression mixes **several** irrational numbers with addition, subtraction, multiplication, or exponents? **Here's how you handle it:** replace each irrational number with its decimal approximation, then carry out the arithmetic step by step. But here's the real secret that saves time and keeps your answer exact—**simplify before you substitute!** For example, $(\sqrt{3})^2 = 3$ exactly (no decimal needed!), and $\sqrt{2} \cdot \sqrt{8} = \sqrt{16} = 4$—the irrational parts cancel out completely. Only use decimal approximations for the parts that *truly* stay irrational. One more important rule: **round at the very end, not in the middle.** Early rounding piles up error and throws off your final answer. The reference line below collects the key approximations you'll need most.



Key Concepts & Quick Review

Useful approximations: $\sqrt{2} \approx 1.41$, $\sqrt{3} \approx 1.73$, $\sqrt{5} \approx 2.24$, $\pi \approx 3.14$

Strategy: Substitute ⇒ Simplify ⇒ Round to the required number of decimal places.

Bounding: Computing both a *lower bound* (using smaller approximations) and an *upper bound* (using larger ones) confirms how accurate the estimate is.

Examples

① Estimate π^2 to two decimal places.

Start by substituting: use $\pi \approx 3.14$. Then compute $\pi^2 \approx 3.14 \times 3.14$. **Break it into parts:** $3 \times 3.14 = 9.42$ and $0.14 \times 3.14 = 0.4396$. **Add them:** $9.42 + 0.44 = 9.86$ (rounding 0.4396 to 0.44). **Check your work:** the actual value is $9.8696\ldots$, so our estimate of 9.86 is spot-on! That's the power of smart substitution.

Answer: $\pi^2 \approx 9.86$

② Estimate $3\sqrt{5} - 2$ to one decimal place.

Scan Me