6 Full-Length OKLAHOMA ALGEBRA II PRACTICE TESTS



Clear practice, student-friendly explanations, and a steady path through Algebra 2.

6 FULL-LENGTH TESTS
Realistic practice to build confidence

STUDENT-FRIENDLY EXPLANATIONS
Clear, step-by-step solutions

BUILD SKILLS
Strengthen concepts and problem-solving

A STEADY PATH
Focused practice to help you reach your goal

SIX-TEST ALGEBRA II WORKBOOK

Oklahoma Algebra II Practice Tests

Clear practice, student-friendly explanations, and a steady path through Algebra 2.

6

FULL-LENGTH TESTS

ALGEBRA II PRACTICE PATH

- Plan
- Practice
- Review
- Grow

1 Start — 2 Build — 3 Check — 4 Improve

Six complete tests
Full-length sets for pacing and stamina.

Warm explanations
Step-by-step answer support after every test.

Oklahoma aligned
Standards references placed before the tests.

Ready to use
Organized for review, tutoring, and independent study.

Practice with purpose. Review with care. Build Algebra II confidence.

Reza Nazari

Copyrighted Material

Copyright ©

Effortless Math Education Inc

Published by Effortless Math Education

www.effortlessmath.com

All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the author, except in the case of brief quotations embodied in critical reviews and certain other noncommercial uses permitted by copyright law, including Section 107 or 108 of the 1976 United States Copyright Act.

This publication is independently produced and has no official connection to any state, district, or national testing program.

Test names and organizational names used herein are the property of their respective trademark holders.

Copyright ©

Copyrighted Material


Pacing rule

Make a first pass through the test for questions you can solve cleanly. Mark the heavy ones, then return with the time you protected. Good pacing is not rushing; it is choosing where your attention goes.

Final check before moving on: Did you answer the exact thing asked? Algebra II problems often ask for one feature of a larger situation. Give the requested feature, not extra work.

Before You Begin Test 1

Let the first test be useful, not perfect



You do not need to feel completely ready before you start. The purpose of a practice test is to show you what is already strong and what still needs attention.

As you work, pay attention to your habits. Do you read graphs carefully? Do you check restrictions? Do you keep signs and exponents organized? Do you return to difficult questions with a calmer mind? These habits are part of the score, even though they do not look like formulas.

Six tests give you room to grow. Use the first one to learn where to begin.

Work carefully. Review honestly. Let each test make the next one stronger.

1. FUNCTIONS, TRANSFORMATIONS, AND INVERSES

Composition: $(f \circ g)(x) = f(g(x))$

Inverse relationship: $f(g(x)) = x$ and $g(f(x)) = x$

Vertical shift: $f(x) + k$ moves up/down

Horizontal shift: $f(x - h)$ moves right by h

Average rate of change: $\dfrac{f(b) - f(a)}{b - a}$

Domain: allowed inputs

Range: possible outputs

2. QUADRATICS, POLYNOMIALS, AND COMPLEX NUMBERS

Quadratic formula: $x = \dfrac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

Vertex: $x = -\dfrac{b}{2a}$

Discriminant: $b^2 - 4ac$

Remainder theorem: remainder is $f(a)$ after division by $x - a$

Complex numbers: $i^2 = -1$

Conjugates: $(a + bi)(a - bi) = a^2 + b^2$

3. EXPONENTIALS, LOGARITHMS, AND SEQUENCES

Exponential model: $y = ab^x$

Log definition: $\log_b a = c \Leftrightarrow b^c = a$

Change of base: $\log_b a = \dfrac{\log a}{\log b}$

Arithmetic: $a_n = a_1 + (n - 1)d$

Geometric: $a_n = a_1 r^{n-1}$

Finite geometric sum: $S_n = a_1 \dfrac{1 - r^n}{1 - r}$

4. TRIGONOMETRY, CONICS, AND STATISTICS

Identity: $\sin^2\theta + \cos^2\theta = 1$

Law of sines: $\dfrac{a}{\sin A} = \dfrac{b}{\sin B}$

Law of cosines: $c^2 = a^2 + b^2 - 2ab\cos C$

Circle: $(x - h)^2 + (y - k)^2 = r^2$

Mean: $\bar{x} = \dfrac{\sum x_i}{n}$

Standard deviation: spread around the mean

Normal curve: about $68\%, 95\%, 99.7\%$

Scan Me

Algebra II

Practice Test 1

1

Questions: 50 **Duration:** No time limit

Calculator Policy: You may use a scientific calculator.

Important Test Information

Question Types:

- **Multiple Choice:** Choose the best answer from the four options (A, B, C, or D).
- **Open-Ended:** Write your complete answer in the space provided.
- **Select All That Apply:** Choose all answers that correctly answer the question.

Test-Taking Tips:

- Read each question carefully before answering.
- Check your answers before moving on.
- Use the formula sheet as needed.

Remember: Take your time, stay calm, and do your best. You've prepared well for this test—you've got this!

Algebra II Reference Materials

QUADRATICS AND POLYNOMIALS

Quadratic formula	$x = \dfrac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
Discriminant	$\Delta = b^2 - 4ac$
Vertex of $ax^2 + bx + c$	$x = -\dfrac{b}{2a}$
Remainder theorem	$f(a)$ is the remainder after division by $x - a$
Difference of squares	$a^2 - b^2 = (a - b)(a + b)$

EXPONENTIALS AND LOGARITHMS

Product rule	$\log_b(MN) = \log_b M + \log_b N$
Quotient rule	$\log_b(M/N) = \log_b M - \log_b N$
Power rule	$\log_b(M^p) = p \log_b M$
Change of base	$\log_b a = \dfrac{\log a}{\log b}$
Compound interest	$A = P(1 + r)^t$

SEQUENCES, SERIES, AND COMPLEX NUMBERS

Arithmetic sequence	$a_n = a_1 + (n - 1)d$
Arithmetic series	$S_n = \dfrac{n}{2}(a_1 + a_n)$
Geometric sequence	$a_n = a_1 r^{n-1}$
Geometric series	$S_n = a_1 \dfrac{1 - r^n}{1 - r}$
Imaginary unit	$i^2 = -1$

TRIGONOMETRY AND CONICS

Pythagorean identity	$\sin^2\theta + \cos^2\theta = 1$
Law of sines	$\dfrac{a}{\sin A} = \dfrac{b}{\sin B} = \dfrac{c}{\sin C}$
Law of cosines	$c^2 = a^2 + b^2 - 2ab\cos C$
Circle	$(x - h)^2 + (y - k)^2 = r^2$
Parabola	$(x - h)^2 = 4p(y - k)$ or $(y - k)^2 = 4p(x - h)$

STATISTICS

Mean	$\bar{x} = \dfrac{\sum x_i}{n}$
Standard deviation idea	typical distance from the mean
Margin of error	critical value $\times$ standard error
Empirical rule	about $68\%, 95\%, 99.7\%$ within $1, 2, 3$ SDs


1) For $x \neq 0$, which expression is equivalent to $x^3 \cdot x^7$?

☐ A. $2x^{10}$
☐ B. x^4
☐ C. x^{10}
☐ D. x^{21}

2) Write $\sqrt[3]{x^2}$ in rational-exponent form, assuming $x \geq 0$.

☐ A. $x^{\frac{3}{2}}$
☐ B. x^6
☐ C. $x^{\frac{2}{3}}$
☐ D. $x^{\frac{1}{6}}$

3) Which point represents the solution to the system of equations shown in the graph?



☐ A. $(0, -3)$
☐ B. $(3, 5)$
☐ C. $(-1, 2)$
☐ D. $(2, 1)$